

# CLASSICAL ELECTRODYNAMICS II

## Homework Set 1

September 9, 2016

1. Consider transverse electromagnetic waves in free space of the form:

$$\begin{aligned}\mathbf{E} &= \mathbf{E}_0(x, y) e^{i(kz - \omega t)}, \\ \mathbf{B} &= \mathbf{B}_0(x, y) e^{i(kz - \omega t)}.\end{aligned}$$

Find the relation between  $k$  and  $\omega$ , as well as the relation between  $\mathbf{E}_0$  and  $\mathbf{B}_0$ . Also show that  $\mathbf{E}_0$  and  $\mathbf{B}_0$  satisfy the equations for electrostatics and magnetostatics in free space; *i.e.*, show that  $\mathbf{E}_0$  and  $\mathbf{B}_0$  have no divergence and no curl.

2. Consider the one-dimensional wave equation,

$$\frac{\partial^2 u}{\partial x^2} - \frac{1}{v^2} \frac{\partial^2 u}{\partial t^2} = 0.$$

Show that by making the substitutions,  $\xi = x - vt$  and  $\eta = x + vt$ , the wave equation takes on a very simple form that can be easily integrated to give the solution,  $u(\xi, \eta) = f(\xi) + g(\eta)$ .

3. Consider a source-free medium in which there is no external magnetization and the polarization vector  $\mathbf{P}$  is a given function of position and time:  $\mathbf{P} = \mathbf{P}(\mathbf{r}, t)$ . Show that the electric field  $\mathbf{E}$  and magnetic induction  $\mathbf{B}$  can be written in terms of a single vector  $\mathbf{Z}$  (called the Hertz vector) as

$$\mathbf{E} = \nabla \times (\nabla \times \mathbf{Z}) - \frac{\mathbf{P}}{\epsilon}, \quad \mathbf{B} = \mu\epsilon \nabla \times \frac{\partial \mathbf{Z}}{\partial t},$$

where with suitable choice of gauge,

$$\nabla^2 \mathbf{Z} - \mu\epsilon \frac{\partial^2 \mathbf{Z}}{\partial t^2} = -\frac{\mathbf{P}}{\epsilon}.$$

How are the usual vector and scalar potentials ( $\mathbf{A}$  and  $\Phi$ ) related to  $\mathbf{Z}$ ?