

CLASSICAL ELECTRODYNAMICS II

Homework Set 5

March 16, 2018

1. Derive a simple general expression for the vector potential in the radiation zone starting with

$$\mathbf{A}(\mathbf{r}, t) = \frac{\mu_0}{4\pi} \int \int G^{(+)}(\mathbf{r}, t; \mathbf{r}', t') \mathbf{J}(\mathbf{r}', t') d^3r' dt' ,$$

where $G^{(+)}(\mathbf{r}, t; \mathbf{r}', t')$ is the retarded (causal) Green function. That is, do *not* assume $\mathbf{J}(\mathbf{r}, t) = \mathbf{J}(\mathbf{r}) e^{-i\omega t}$.

2. Now consider a rotating electric dipole consisting of two equal and opposite charges q and $-q$ attached to the ends of a rod of length s . The rod rotates counterclockwise in the x - y plane with angular speed $\omega = ck$. The electric dipole moment of the system at $t = 0$ has the value $\mathbf{p}_0 = qs\hat{x}$. Use your result from part (a) to calculate $\mathbf{A}(\mathbf{r}, t)$ in the radiation zone. (*Hint*: Recall that $ks \ll 1$.) Show that your result can be put in the complex form:

$$\mathbf{A}(\mathbf{r}, t) = \frac{\mu_0}{4\pi r} p_0 \omega \left(\hat{\phi} - i\hat{\rho} \right) e^{i(\phi - \omega t + kr)} ,$$

where $\hat{\rho} = \hat{x} \cos \phi + \hat{y} \sin \phi$ and $\hat{\phi} = -\hat{x} \sin \phi + \hat{y} \cos \phi$.

3. Use the complex expression in part (b) to calculate $\mathbf{B}(\mathbf{r}, t)$ in the radiation zone.
4. Use your result of part (c) to calculate $\mathbf{E}(\mathbf{r}, t)$ in the radiation zone.
5. Use your results of parts (c) and (d) to determine equations for the real, physical fields \mathbf{E} and \mathbf{B} . Then calculate the instantaneous Poynting vector. Is the instantaneous Poynting vector azimuthally symmetric (independent of ϕ)? If not, are there any observation points where it is independent of ϕ ? Finally, calculate the time-averaged Poynting vector and comment on whether it depends on ϕ or not.
6. Lastly calculate the time-averaged power radiated per unit solid angle and make a sketch of the radiation pattern.