

# Measurements of open charm production and flow in 200 GeV Au+Au collisions with the STAR experiment at RHIC


*Spiros Margetis for the STAR Collaboration  
Kent State University*



# Introduction

Large collective flow and suppression of yields for charm hadrons in 200 GeV A+A collisions have been already reported by STAR

**New data:** Understand better heavy quark production, transport and hadronization in the presence of QGP




**New [high statistics/optimized] extensive measurements by STAR!**

- Large **directed\*** ( $v_1$ ) and **elliptic** ( $v_2$ ) flow of  $D^0$
- Hadronization:  $\Lambda_c$ ,  $D_s$
- **In medium energy loss:  $D^0$ , B-mesons\***
- Medium modifications to yields/life-time:  $D^{*+/-}$
- **Total charm cross-section**


\* Not reported here. See QM2018 talks of S. Sinha and S. Radhakrishnan

# The STAR Detector



- 2 layers of Si pixels with MAPS and 2 layers of Si strips
- Full azimuthal coverage



*Phys. Rev. Lett. 118 (2017) 212301*



STAR Heavy Flavor Tracker (HFT) provides excellent vertex/track-dca resolution and allows reconstruction of charm hadron decays

# Recent $D^0$ Elliptic Flow ( $v_2$ ) Results from STAR

L Adamczyk et. al. (STAR Collaboration),  
Phys Rev. Lett. 118, 212301 (2017)





- STAR published  $D^0$   $v_2$  from data taken during 2014 run
- $D^0$  flow magnitude consistent with NCQ scaling in minimum bias and mid-central collisions.
- High statistics 2016 run data allow to improve precision of the charm flow measurements at RHIC energy
- The 2016 data also allow us to extend NCQ scaling test to finer centrality bins

Precise  $D^0$   $v_2$  measurement can allow:

➔ Quantitative studies of QGP properties (transport coefficients)





# D<sup>0</sup> v<sub>2</sub> Comparison to Light Hadrons



- D<sup>0</sup> v<sub>2</sub> results from combined 2014 + 2016 data
- D<sup>0</sup> v<sub>2</sub> measurement extended to 0-10% centrality
- Clear mass ordering for  $p_T < 2$  GeV/c in 10-40% centrality
- D<sup>0</sup> v<sub>2</sub> for  $p_T > 2$  GeV/c in 10-40% centrality follows the mesons

# NCQ Scaling Test




- NCQ scaling test with improved precision in  $D^0$   $v_2$  measurement
- NCQ-scaled  $D^0$   $v_2$  consistent with light hadrons for  $(m_T - m_0) / n_q < 2.5$   $\text{GeV}/c^2$  in 10-40%
- Evidence of charm quarks flowing with the medium

Charm quark appear to have achieved thermal equilibrium with the medium



# D<sup>0</sup> v<sub>2</sub>: Data vs. Models



Compared Models	x <sup>2</sup> /NDF	p-value
SUBATECH [1]	17.3/8	0.026
TAMU c quark diff. [2]	12.0/8	0.15
TAMU no c quark diff. [2]	33.7/8	4.5 x 10 <sup>-5</sup>
Duke (Bayesian) [3]	8.5/8	0.39
3D viscous hydro [4]	3.7/6	0.71
LBT [5]	13.3/8	0.10
PHSD [6]	8.7/7	0.27
Catania [7]	9.7/8	0.29

- [1] SUBATECH: *Phys Rev C* 90, 054909 (2014), *Phys Rev C* 92, 014910 (2015)  
 [2] TAMU: *Phys Rev C* 86, 014903 (2012), *Phys Rev Lett* 110, 112301 (2013)  
 [3] Duke: *Phys Rev C* 92, 024907 (2015)  
 [4] 3D viscous hydro: *Phys Rev C* 86, 024911 (2012)  
 [5] LBT: *Phys Rev C* 94, 014909 (2016)  
 [6] PHSD: *Phys ReV* 90, 051901 (2014), *Phys ReV* 90, 051901 (2014)  
 [7] Catania: *Phys ReV* 96, 044905 (2017)


- D<sup>0</sup> v<sub>2</sub> results from combined 2014 + 2016 data
- Improved precision to constrain the models




# $\Lambda_c$ and Heavy Flavor Hadronization

- Strong enhancement of  $\Lambda_c/D^0$  ratio seen in Au+Au collisions by STAR
- Enhancement predicted from coalescence hadronization
- An enhancement relative to PYTHIA also seen in p+p and p+Pb collisions at LHC

ALICE: arXiv:1712.09581



Ko: PRC 79 (2009) 044905  
 Greco: PRD 90 (2014) 054018  
 SHM: PRC 79 (2009) 044905




- How does  $\Lambda_c$  production change from peripheral to central A+A collisions?
- What is the  $p_T$  dependence of  $\Lambda_c$  production in A+A collisions?





# $p_T$ Dependence of $\Lambda_c/D^0$ Ratio




Ko: *Phys.Rev.C* 79 (2009) 044905  
 Greco: *Eur.Phys.J.C* (2018) 78:348  
 SHM: *Phys.Rev.C* 79 (2009) 044905

- Strong enhancement of  $\Lambda_c$  production compared to PYTHIA calculations
- Enhancement increases towards low  $p_T$
- Coalescence model predictions are closer to data, but the observed enhancement is larger than that predicted by models, particularly at higher  $p_T$
- Ratio not described by the Statistical Hadronization Model



# Centrality Dependence of $\Lambda_c$ Production





ALICE: arXiv:1712.09581

- First measurement of centrality dependence of  $\Lambda_c$  production in heavy-ion collisions
- $\Lambda_c/D^0$  ratio increases from peripheral to central, indicative of hot medium effects
- Ratio for peripheral Au+Au consistent with p+p values at 7 TeV



# D<sub>s</sub> Production

- D<sub>s</sub>/D<sup>0</sup> enhancement expected in central A+A collisions, from strangeness enhancement and coalescence hadronization



- D<sub>s</sub> yield (relative to D<sup>0</sup>) is enhanced in A+A collisions
- Enhancement is larger than model predictions, particularly at higher  $p_T$
- Ratio close to SHM predictions

*ep/pp/ep avg: M Lisovsky, et. al. EPJ C 76, 397 (2016)*


*TAMU: H. Min et al. PRL 110, 112301 (2013)*

*SHM: A. Andronic et al., PLB 571 (2003) 36*



# D<sup>0</sup> Spectra and R<sub>AA</sub>



- Updated results from STAR for D<sup>0</sup> extending to low p<sub>T</sub> and non-central collisions



- $R_{AA}$  in central events  $< 1$  at all  $p_T$
- Suppression at high  $p_T$  increases with centrality




# D<sup>0</sup> Cross-section and BW Fits to Spectra





- Total  $D^0$  cross-section is nearly independent of centrality, and smaller than in p+p. However, decreases towards central collisions for  $p_T > 4$  GeV/c
- Blast Wave fits to  $D^0$  spectra:
  - BW fits to  $p_T < 5$  GeV/c. Both standard and Tsallis BW fits tried
  - Result suggest an earlier freeze-out for  $D^0$  than light hadrons

# D\* Production in Au+Au Collisions

- Measure  $D^{*+}/D^0$  ratio
- $D^{*+}$  feed-down contribution to  $D^0$  yields (  $D^{*+} \rightarrow D^0 \pi_{soft}^+$  )
- In-medium effects:
  - Shorter life time in medium (?). Lifetime in vacuum is  $\sim 2000$  fm/c, but spectral function predicted to broaden in medium (*R.Rapp et.al Phys. Rev. C (2018)97, 034918* )
  - Rescattering can lead to loss of yield which was already seen for  $K^*$  (*STAR, Phys. Rev. C (2011)84, 034909*)



# D\* Production in Au+Au Collisions



- D\*<sup>+</sup>/D<sup>0</sup> ratio consistent with PYTHIA and with ALICE data at higher p<sub>T</sub>
- Ratio of the integrated yields shows no strong centrality dependence



# Total Charm Cross-section

- Total charm cross-section is estimated from the various charm hadron measurements

- $D^0$  yields are measured down to zero  $p_T$
- For  $D^{+/-}$  and  $D_s$ , Levy (power law) fits to measured spectra are used for extrapolation (systematics).
- For  $\Lambda_c$ , three model fits to data are used and differences are included in systematics

Charm Hadron		Cross Section $d\sigma/dy$ ( $\mu\text{b}$ )
Au+Au 200 GeV (10-40%)	$D^0$	$41 \pm 1 \pm 5$
	$D^+$	$18 \pm 1 \pm 3$
	$D_s^+$	$15 \pm 1 \pm 5$
	$\Lambda_c^+$	$78 \pm 13 \pm 28^*$
	<b>Total</b>	<b><math>152 \pm 13 \pm 29</math></b>
p+p 200 GeV	<b>Total</b>	<b><math>130 \pm 30 \pm 26</math></b>

\* derived using  $\Lambda_c^+ / D^0$  ratio in 10-80%

- Total charm cross-section is consistent with p+p value within uncertainties.





# Summary

- **Extensive measurements of charm hadron yields in heavy-ion collisions by STAR**
  - Combined 2014+2016 data
  - Improved significance from supervised machine-learning algorithms
- **Large  $D^0$  elliptic flow STAR**
  - Improved precision of  $D^0$   $v_2$  results with combined 2014 and 2016 data
  - $D^0$   $v_2$  result suggests charm quarks achieve a thermal equilibrium with the medium
  - Precise  $D^0$   $v_2$  measurements can further constrain model calculations
- **Strong modification of charm hadron spectra and hadrochemistry in A+A collisions!**
  - Total charm cross-section consistent with p+p within systematic uncertainties.
  - Strong enhancement seen for  $\Lambda_c/D^0$  ratio in Au+Au. Suggests coalescence hadronization of deconfined charm quarks in the medium
  - Strong suppression of  $D^0$  yields at higher  $p_T$  in most central collisions



THANK YOU



Back Up






# Summary


## Directed flow

- First evidence of non-zero directed flow for heavy flavor
- Both  $D^0$  and  $\bar{D}^0$  show negative  $v_1$ -slope near mid-rapidity
- Heavy flavor  $v_1 >$  light flavor  $v_1$   
Data can be used to probe initial matter distribution
- Current precision is not sufficient to draw conclusion on magnetic field induced charge separation of heavy quarks




## Elliptic flow

- Improved precision of  $D^0$   $v_2$  results with combined 2014 and 2016 data
- $D^0$   $v_2$  result suggests charm quarks achieve a thermal equilibrium with the medium
- Precise  $D^0$   $v_2$  measurements can further constrain model calculations



# Boosted Decision Trees (BDT) for $\Lambda_c$ Signal Extraction


- Simple cuts on variables have limitations on signal-background separation
- Supervised learning algorithms can do better!
  - Boosted Decision Trees: successive binary cuts on attributes
  - Good performance for classification problems
  - 7 topological variables as input
  - For training: signal from MC (with detector effects), background from data




# Boosted Decision Trees (BDT) for $\Lambda_c$ Signal Extraction

- Simple cuts on variables have limitations on signal-background separation
- Supervised learning algorithms can do better!

QM18



QM17




- More than 50% improvement in signal significance with TMVA BDT.
- Also new data from 2016  $\rightarrow$  Effectively 4x more data compared to QM17



# Non-prompt $D^0$

- Charm quarks interact strongly with the medium. How about bottom?
- Is there mass hierarchy for energy loss? Is  $\Delta E_c > \Delta E_b$  ?




- $R_{AA}$  of B mesons estimated from the measured non-prompt  $D^0$  fraction
- Need better statistics and improved precision to understand mass dependence of energy loss.


# Non-prompt $D^0$

- Charm quarks interact strongly with the medium. How about bottom?
- Is there mass hierarchy for energy loss? Is  $\Delta E_c > \Delta E_b$  ?


- Improved signal significance for non-prompt  $D^0$  fraction using BDT
- New results with 2014+2016 data on the way



- $R_{AA}$  of B mesons estimated from the measured non-prompt  $D^0$  fraction
- Need better statistics and improved precision to understand mass dependence of energy loss.








Charm quarks seem to acquire the same flow as light quarks!



# Back Up II



# D<sup>+/-</sup> R<sub>AA</sub>





- Similar suppression for D<sup>0</sup> and D<sup>+/-</sup>
- Spectra measurements important for total charm cross-section






# Erratum details

## Erratum: $D^0$ in AuAu (2010/2011 TPC Analysis) - I PRL 113 (2014) 142301

- Two mistakes were discovered in calculating TOF related efficiency corrections
  - Hybrid PID: algorithm inconsistently implemented in data analysis vs efficiency calculation
  - a transverse distance of closest approach cut efficiency was included in the correction two times
- p+p measurement: no issue ( $D^0$  at  $p_T < 2$  GeV/c +  $D^*$  at 2-6 GeV/c, *PRD 86 (2012) 072012*), but the p+p  $D^0$  baseline used for  $R_{AA}$  is updated with latest knowledge of charm frag. ratios
  - considering the  $p_T$  dependence of  $D^*/D^0$  frag. ratio
  - latest world average of  $c \rightarrow D^0$  and  $c \rightarrow D^*$  frag. ratios





# Total charm cross-section: procedure



- Extracted for 10-40% centrality.
- Yields for  $D^{+/-}$  and  $\Lambda_c$  are scaled to 10-40% centrality using measured ratio to  $D^0$ .
- Uncertainty evaluation and propagation:
  - In the  $p_T$  range with data points:
    - point by point statistical error propagated
    - point by point systematic error propagated
  - In the  $p_T$  range without data points
    - uncertainties from fit to points with statistical + systematic error
    - extrapolation uncertainty from variation of fit function

# BDT vs Rectangular Cuts Comparison


- Simple cuts on variables have limitations on signal-background separation
- Supervised learning algorithms can do better!



- More than 50% improvement in signal significance with TMVA BDT.




# BW fits to $D^0$ spectra



- Fit values shown were from BW fits
- TBW gives lower temperatures for all particles, but similar radial flow



# $R_{AA}$ of B through different channels



- The decay kinematics need to be unfolded for a fair comparison among different channels.

